

QGP Physics – from Fixed Target to LHC

2. Kinematic Variables

Prof. Dr. Johanna Stachel, PD Dr. Klaus Reygers
Physikalisches Institut
Universität Heidelberg
SS 2011

Notations and Conventions

Natural units: $c = \hbar = 1$ Also: $k_B = 1 \rightarrow E = k_B T$, $T = 2 \cdot 10^{12} \text{ K} \approx 172 \text{ MeV}$

Space-time coordinates
(contravariant vector): $x^\nu = (x^0, x^1, x^2, x^3) = (t, \vec{x}) = (t, x, y, z)$

Relativistic energy and momentum: $E = \gamma m$, $p = \gamma \beta m$, $m = \text{rest mass}$


4-momentum vector: $p^\mu = (p^0, p^1, p^2, p^3) = (E, \vec{p}) = (E, \vec{p}_T, p_z) = (E, p_x, p_y, p_z)$

Scalar product of two 4-vectors a and b : $a \cdot b = a^0 b^0 - \vec{a} \cdot \vec{b}$

Relation between energy and momentum: $E^2 = p^2 + m^2$

Center-of-Momentum System (CMS)

Consider a collision of two particles. The CMS is defined by $\vec{p}_a = -\vec{p}_b$


$$p_a = (E_a, \vec{p}_a) \quad p_b = (E_b, \vec{p}_b)$$


The Mandelstam variable s is defined as $s := (p_a + p_b)^2 \stackrel{CMS}{=} (E_a + E_b)^2$

The center-of-mass energy \sqrt{s} is the total energy available in the CMS


\sqrt{s} for Fixed-Target und Collider Experiments

Fixed-target experiment:



$$\sqrt{s} = \begin{cases} \sqrt{m_1^2 + m_2^2 + 2E_1^{\text{lab}}m_2} & E_1^{\text{lab}} \gg m_1, m_2 \\ \sqrt{2E_1^{\text{lab}}m_2} & \approx \end{cases}$$

Collider:



$$\sqrt{s} = \sqrt{m_1^2 + m_2^2 + 2E_1^{\text{lab}}E_2^{\text{lab}} + 2p_1^{\text{lab}}p_2^{\text{lab}}} \\ \vec{p}_1 = -\vec{p}_2 \\ m_1 = m_2 \\ 2E_1^{\text{lab}}$$

Example: Anti proton production

(fixed-target experiment): $p + p \rightarrow p + p + p + \bar{p}$

Minimum energy required to produce an anti-proton:


In CMS, all particles at rest after the reaction, i.e., $\sqrt{s} = 4 m_p$, hence:

$$E_1^{\text{lab}, \min} = \frac{(4m_p)^2 - 2m_p^2}{2m_p} = 7m_p$$

Rapidity

The rapidity y is a generalization of velocity $\beta_L = p_L/E$:

$$y := \operatorname{arctanh} \beta_L = \frac{1}{2} \ln \frac{1 + \beta_L}{1 - \beta_L} = \frac{1}{2} \ln \frac{E + p_L}{E - p_L}$$




For small velocities: $y \approx \beta_L$ for $\beta_L \ll 1$

With $e^y = \sqrt{\frac{E + p_L}{E - p_L}}$, $e^{-y} = \sqrt{\frac{E - p_L}{E + p_L}}$

and $\sinh x = \frac{1}{2} (e^x - e^{-x})$, $\cosh x = \frac{1}{2} (e^x + e^{-x})$

we readily obtain $E = m_T \cdot \cosh y$, $p_L = m_T \cdot \sinh y$

where $m_T := \sqrt{m^2 + p_T^2}$ is called the *transverse mass*



Rapidity II

y is not Lorentz invariant, however, it has a simple transformation property:

$$y = y' + y_{S'}$$

↑
rapidity in S'

Rapidity in system S

Rapidity of S' measured in S , defined as
 $y_{S'} = \frac{1}{2} \ln \frac{1 + \beta_{S'}}{1 - \beta_{S'}}$

Consider collisions of two particles with equal mass m and rapidities y_a and y_b .
The rapidity of the CMS y_{CM} is then given by:

$$\bullet \xrightarrow{m, y_a} \quad \xleftarrow{m, y_b} \bullet \qquad y_{CM} = (y_a + y_b)/2$$

In the center-of-mass frame, the rapidities of a and b are:

$$y_a^* = -(y_b - y_a)/2 \qquad \text{and} \qquad y_b^* = (y_b - y_a)/2$$

Pseudorapidity

$$y = \frac{1}{2} \ln \frac{E + p \cos \vartheta}{E - p \cos \vartheta} \stackrel{p \gg m}{\approx} \frac{1}{2} \ln \frac{1 + \cos \vartheta}{1 - \cos \vartheta} = \frac{1}{2} \ln \frac{2 \cos^2 \frac{\vartheta}{2}}{2 \sin^2 \frac{\vartheta}{2}} = -\ln \left[\tan \frac{\vartheta}{2} \right] =: \eta$$

$\cos(2\alpha) = 2 \cos^2 \alpha - 1 = 1 - 2 \sin^2 \alpha$

Especially: $y = \eta$ for $m = 0$

Analogous to the relations for the rapidity we find

$$p = p_T \cdot \cosh \eta, \quad p_L = p_T \cdot \sinh \eta$$

Example: Beam Rapidities

$$y = \frac{1}{2} \ln \frac{E + p_z}{E - p_z} = \ln \frac{E + p_z}{\sqrt{E^2 - p_z^2}} = \ln \frac{E + p_z}{m} \approx \ln \frac{2E}{m}$$

Beam momentum (GeV/c)	Beam rapidity
100	5.36
158	5.81
2750	8.86
3500	8.92
7000	9.61

Quick Overview: Kinematic Variables

Transverse momentum


$$p_T = p \sin \vartheta$$

rapidity


$$y = \text{atanh } \beta_L$$

pseudorapidity

$$\eta = -\ln \tan \frac{\vartheta}{2}$$



Example of a Pseudo-rapidity Distribution



Beam rapidity:

$$y_{beam} = \ln \frac{E + p}{m} = 5.36$$

Average number of charged particles per collision:


$$\langle N_{ch} \rangle = \int \frac{dN_{ch}}{d\eta} d\eta \approx 20$$

Difference between dN/dy and $dN/d\eta$ in the CMS





$$\frac{dN}{d\eta} = \sqrt{1 - \frac{m^2}{m_T^2 \cosh^2 y}} \frac{dN}{dy}$$

Difference between dN/dy and $dN/d\eta$ in the CMS at $y = 0$:


$$\frac{dN/d\eta}{dN/dy} \approx \sqrt{1 - \frac{m^2}{\langle m_T^2 \rangle}}$$




Pseudorapidity Distribution and Angular Distribution



Total Energy and Transverse Energy



$$\frac{dE}{dy} \quad E = \sqrt{m^2 + p^2}$$



$$\frac{dm_T}{dy} \quad m_T = \sqrt{m^2 + p_T^2}$$

Luminosity and Cross Sections (I)

The luminosity L of a collider is defined by:

$$N_{\text{int}} = \sigma \cdot L$$

L = luminosity(in $\text{s}^{-1}\text{cm}^{-2}$)

N_{int} = Number of interactions of a certain type per second

σ = cross section for this reaction

$$L = \frac{n_1 n_2 f}{A}$$

n_1, n_2 = numbers of particles per bunch in the two beams

f = bunch crossing frequency at a given crossing point

A = beam crossing area

Luminosity and Cross Sections (II)

The luminosity can be determined by measuring the beam current:

$$I_{1,2} = n_{1,2} \cdot N_b \cdot e \cdot f$$

N_b = number of bunches in the beam

e = elementary electric charge

The crossing area A is usually calculated as

$$A = 4\pi\sigma_x\sigma_y$$

The standard deviations of the beam profiles are measured by sweeping the beams transversely across each other in a so called van der Meer scan.

Integrated luminosity:

$$L_{\text{int}} = \int L \, dt$$

Lorentz invariant Phase Space Element

If one is interested in the production of a particle X one could define the observable

$$\frac{1}{L_{\text{int}}} \frac{d^3 N_x}{d\vec{p}} = \frac{1}{L_{\text{int}}} \frac{d^3 N_x}{dp_x dp_y dp_z}$$

However, the phase space density would then not be Lorentz invariant

$$\frac{d^3 N}{dp'_x dp'_y dp'_z} = \frac{\partial(p_x, p_y, p_z)}{\partial(p'_x, p'_y, p'_z)} \cdot \frac{d^3 N}{dp_x dp_y dp_z} = \frac{E}{E'} \cdot \frac{d^3 N}{dp_x dp_y dp_z}$$

We thus use the Lorentz invariant phase space element $\frac{d^3 \vec{p}}{E} = \frac{dp_x dp_y dp_z}{E}$

The corresponding observable is called Lorentz invariant cross section:

$$E \frac{d^3 \sigma}{d^3 \vec{p}} = \frac{1}{L_{\text{int}}} E \frac{d^3 N_x}{d^3 \vec{p}} = \underbrace{\frac{1}{N_{\text{evt,tot}}} E \frac{d^3 N_x}{d^3 \vec{p}}}_{\text{this is called the invariant yield}} \sigma_{\text{tot}}$$

[Lorentz invariant Phase Space Element: Proof of Invariance]

Lorentz boost along the z axis:

$$\begin{aligned} p'_x &= p_x \\ p'_y &= p_y \\ p'_z &= \gamma(p_z - \beta E), & p_z = \gamma(p'_z + \beta E') \\ E' &= \gamma(E - \beta p_z), & E = \gamma(E' + \beta p'_z) \end{aligned}$$

Jacobian:

$$\frac{\partial(p_x, p_y, p_z)}{\partial(p'_x, p'_y, p'_z)} = \begin{vmatrix} \frac{\partial p_x}{\partial p'_x} & 0 & 0 \\ 0 & \frac{\partial p_y}{\partial p'_y} & 0 \\ 0 & 0 & \frac{\partial p_z}{\partial p'_z} \end{vmatrix}$$

$$\frac{\partial p_x}{\partial p'_x} = 1, \quad \frac{\partial p_y}{\partial p'_y} = 1, \quad \frac{\partial p_z}{\partial p'_z} = \frac{\partial}{\partial p'_z} [\gamma(p'_z + \beta E')] = \gamma \left(1 + \beta \frac{\partial E'}{\partial p'_z} \right)$$

$$\frac{\partial E'}{\partial p'_z} = \frac{\partial}{\partial p'_z} \left[(m^2 + p'^2_x + p'^2_y + p'^2_z)^{1/2} \right] = \frac{p'_z}{E'}$$

$$\rightsquigarrow \frac{\partial p_z}{\partial p'_z} = \gamma \left(1 + \beta \frac{p'_z}{E'} \right) = \frac{E}{E'}$$

And so we finally obtain:

$$\frac{\partial(p_x, p_y, p_z)}{\partial(p'_x, p'_y, p'_z)} = \frac{E}{E'}$$

Invariant Cross Section

Invariant cross section in practice:

$$\begin{aligned}
 E \frac{d^3\sigma}{d^3p} &= E \frac{1}{p_T} \frac{d^3\sigma}{dp_T dp_z d\varphi} \\
 dp_z/dy = m_T &\equiv \cosh y = E \quad \frac{1}{p_T} \frac{d^3\sigma}{dp_T dy d\varphi} \\
 \text{symmetry in } \varphi &\equiv \frac{1}{2\pi p_T} \frac{d^2\sigma}{dp_T dy}
 \end{aligned}$$

Sometimes also measured as a function of m_T :

$$\frac{1}{2\pi m_T} \frac{d^2\sigma}{dm_T dy} = \frac{1}{2\pi m_T} \frac{d^2\sigma}{dp_T dy} \frac{dp_T}{dm_T} = \frac{1}{2\pi p_T} \frac{d^2\sigma}{dp_T dy}$$


Integral of the inv. cross section:

$$\int E \frac{d^3\sigma}{d^3p} d^3p/E = \langle N_x \rangle \cdot \sigma_{\text{tot}}$$

/

Average yield of particle X
per event

Example: Invariant cross section for neutral pion production in p+p at $\sqrt{s} = 200$ GeV




Invariant Mass

Consider the decay of a particle in two daughter particles. The mass of the mother particle is given by (“invariant mass”):


$$\begin{aligned}
 M^2 &= \left[\left(\frac{E_1}{\vec{p}_1} \right) + \left(\frac{E_2}{\vec{p}_2} \right) \right]^2 = (E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2 \\
 &= m_1^2 + m_2^2 + 2E_1 E_2 - 2\vec{p}_1 \cdot \vec{p}_2 \\
 &= m_1^2 + m_2^2 + 2E_1 E_2 - 2p_1 p_2 \cos \vartheta
 \end{aligned}$$

Example: π^0 decay: $\pi^0 \rightarrow \gamma + \gamma$, $m_1 = m_2 = 0$, $E_i = p_i$

$$\Rightarrow M = \sqrt{2E_1 E_2 (1 - \cos \vartheta)}$$



Example:
 π^0 peak with
combinatorial
background



Points to Take Home

Center-of-mass energy \sqrt{s} : Total energy in the center-of-mass (or momentum) system (rest mass of + kinetic energy)

Observables: Transverse momentum p_T and rapidity y

Pseudorapidity $\eta \approx y$ for $E \gg m$ ($\eta = y$ for $m = 0$, e.g., for photons)

Production rates of particles described by the Lorentz invariant cross section:

Lorentz-invariant cross section: $E \frac{d^3\sigma}{d^3p}$